arXiv:1412.3535v1 [math.FA] 11 Dec 2014

ALMOST COMMUTING FUNCTIONS OF ALMOST COMMUTING SELF-ADJOINT OPERATORS

Aleksei Aleksandrov[a], Vladimir Peller[b]

[a] *St-Petersburg Branch, Steklov Institute of Mathematics, Fontanka 27, 191023 St-Petersburg, Russia and Department of Mathematics and Mechanics, Saint Petersburg State University, 28, Universitetski pr., St.Petersburg, 198504, Russia*
[b] *Department of Mathematics, Michigan State University, East Lansing, MI 48824, USA*

Abstract. Let A and B be almost commuting (i.e, $AB-BA\in\boldsymbol{S}_1$) self-adjoint operators. We construct a functional calculus $\varphi\mapsto\varphi(A,B)$ for φ in the Besov class $B^1_{\infty,1}(\mathbb{R}^2)$. This functional calculus is linear, the operators $\varphi(A,B)$ and $\psi(A,B)$ almost commute for φ, $\psi\in B^1_{\infty,1}(\mathbb{R}^2)$, $\varphi(A,B)=u(A)v(B)$ whenever $\varphi(s,t)=u(s)v(t)$, and the Helton–Howe trace formula holds. The main tool is triple operator integrals.

Fonctions presque commutant d'opérateurs auto-adjoints presque commutant

Résumé. On dit que des opérateurs A et B presque commutent si leur commutateur $[A,B]$ appartient à classe trace. Pour des opérateurs A et B auto-adjoints qui presque commutent nous construisons un calcul fonctionnel $\varphi\mapsto\varphi(A,B)$, $\varphi\in B^1_{\infty,1}(\mathbb{R}^2)$, où $B^1_{\infty,1}(\mathbb{R}^2)$ est la classe de Besov. Ce calcul a des propriétés suivantes: il est linéaire, les opérateurs $\varphi(A,B)$ et $\psi(A,B)$ presque commutent pour toutes les fonctions φ et ψ dans $B^1_{\infty,1}(\mathbb{R}^2)$, $\varphi(A,B)=u(A)v(B)$ si $\varphi(s,t)=u(s)v(t)$, et la formule des traces de Helton et Howe est vraie. L'outil principal est les intégrales triples opératorielles.

Version française abrégée

Soient A et B des opérateurs auto-adjoints qui *presque commutent*, c'est-à-dire leur commutateur $[A,B]\stackrel{\rm def}{=}AB-BA$ appartient à classe trace $\boldsymbol{S}_1$. Dans [HH] on a obtenu la formule suivante:

$$\operatorname{trace}\big(\mathrm{i}\big(\varphi(A,B)\psi(A,B)-\psi(A,B)\varphi(A,B)\big)\big)=\iint_{\mathbb{R}^2}\left(\frac{\partial\varphi}{\partial x}\frac{\partial\psi}{\partial y}-\frac{\partial\varphi}{\partial y}\frac{\partial\psi}{\partial x}\right)dP$$

pour tous les polynômes φ et ψ de deux variables, où P est une mesure signée boreliienne à support compact. Il s'est trouvé que la mesure P est absolument continue et

$$dP(x,y)=\frac{1}{2\pi}g(x,y)\,dx\,dy,$$

où g est la fonction principale de Pincus (voir la partie anglaise pour plus d'informations).

Le calcul polynomial $\varphi\mapsto\varphi(A,B)$ était étendu dans [CP] et [Pe4]. Dans [Pe4] on a construit le calcul fonctionnel $\varphi\mapsto\varphi(A,B)$ pour φ appartenant à l'intersection des produit tensoriels projectifs $\mathcal{C}\stackrel{\rm def}{=}\big(L^\infty(\mathbb{R})\hat{\otimes}B^1_{\infty,1}(\mathbb{R})\big)\bigcap\big(B^1_{\infty,1}(\mathbb{R})\hat{\otimes}L^\infty(\mathbb{R})\big)$. Ce calcul est linear, les opérateurs $\varphi(A,B)$ et $\psi(A,B)$ presque commute pour φ, $\psi\in\mathcal{C}$. En outre si $\varphi(s,t)=u(s)v(t)$, alors $\varphi(A,B)=u(A)v(B)$. Finalement, la formule de Helton et Howe si-dessus est vrai pour φ et ψ dans $\mathcal{C}$.

Il était aussi démontré dans [Pe4] que c'est impossible de construire un calcul fonctionnel $\varphi\mapsto\varphi(A,B)$ pour toutes les fonctions φ continument dérivables qui ait les propriétés si-dessus.

Le but de cette note est d'améliorer les résultats de [Pe4].

Pour une fonction φ dans la classe de Besov $B^1_{\infty,1}(\mathbb{R}^2)$ nous définissons l'opérateur $\varphi(A,B)$ comme ça:

$$\varphi(A,B)=\iint f(x,y)\,dE_A(x)\,dE_B(y),$$

où E_A et E_B sont les mesures spéctrales des opérateurs A et B. La théorie d'intégrales doubles opératorielles était développée par Birman et Solomyak [BS] (voir aussi [Pe2] et [AP]).

Le résultat principal de cette note est basé sur la formule suivante:

$$\big[\varphi(A,B),Q\big] = \iint\!\!\!\int \frac{\varphi(x,y_1)-\varphi(x,y_2)}{y_1-y_2}\,dE_A(x)\,dE_B(y_1)[B,Q]\,dE_B(y_2)$$
$$+ \iint\!\!\!\int \frac{\varphi(x_1,y)-\varphi(x_2,y)}{x_1-x_2}\,dE_A(x_1)[A,Q]\,dE_A(x_2)\,dE_B(y), \tag{1}$$

où $\varphi \in B^1_{\infty,1}(\mathbb{R}^2)$ et A et B sont des opérateurs auto-adjoints pour lesquels les commutateurs $[A,Q]$ et $[B,Q]$ appartiennent à $\boldsymbol{S}_1$.

Les intégrales triples opératorielles

$$\iint\!\!\!\int \Phi(x_1,x_2,x_3)\,dE_1(x_1)T\,dE_2(x_2)R\,dE_3(x_3) \tag{2}$$

étaint définies dans [Pe6] pour les fonctions Φ dans le produit tensoriel projectif intégral; ici T et R sont des opérateurs bornés et E_1, E_2, E_3 sont des mesures spéctrales. Puis dans [JTT] les intégrales triples opératorielles étaint définies pour les fonctions qui appartiennent au produit tensoriel de Haagerup $L^\infty(E_1)\otimes_{\rm h}L^\infty(E_2)\otimes_{\rm h}L^\infty(E_2)$ (voir [Pis]). Il était établi dans [ANP] que les conditions $T\in\boldsymbol{S}_1$ ou $R\in\boldsymbol{S}_1$ et $\Phi\in L^\infty(E_1)\otimes_{\rm h}L^\infty(E_2)\otimes_{\rm h}L^\infty(E_2)$ n'impliquent pas que l'opérateur (2) appartienne à classe trace.

On a défini dans [ANP] les produit tensoriels du type de Haagerup $L^\infty(E_1)\otimes_{\rm h}L^\infty(E_2)\otimes^{\rm h}L^\infty(E_3)$ et $L^\infty(E_1)\otimes^{\rm h}L^\infty(E_2)\otimes_{\rm h}L^\infty(E_3)$ (voir la partie anglaise). On a démontré dans [ANP] que si $T\in\boldsymbol{S}_1$, R est un opérateur borné et $\Phi\in L^\infty(E_1)\otimes_{\rm h}L^\infty(E_2)\otimes^{\rm h}L^\infty(E_3)$, alors l'opérateur (2) appartient à $\boldsymbol{S}_1$. De même, si $R\in\boldsymbol{S}_1$, T est un opérateur borné et $\Phi\in L^\infty(E_1)\otimes^{\rm h}L^\infty(E_2)\otimes_{\rm h}L^\infty(E_3)$, alors l'opérateur (2) appartient à $\boldsymbol{S}_1$.

Si $\varphi\in B^1_{\infty,1}(\mathbb{R}^2)$, alors la fonction $(x_1,x_2,y)\mapsto\big(\varphi(x_1,y)-\varphi(x_2,y)\big)(x_1-x_2)^{-1}$ appartient à l'espace $L^\infty(E_A)\otimes_{\rm h}L^\infty(E_A)\otimes^{\rm h}L^\infty(E_B)$ et la fonction $(x,y_1,y_2)\mapsto\big(\varphi(x,y_1)-\varphi(x,y_2)\big)(y_1-y_2)^{-1}$ appartient à $L^\infty(E_A)\otimes_{\rm h}L^\infty(E_B)\otimes^{\rm h}L^\infty(E_B)$, voir [ANP]. Donc les intégrales dans (3.1) sont bien définis est appartiennent à $\boldsymbol{S}_1$.

La formule (3.1) implique que si φ et ψ appartiennent à $B^1_{\infty,1}(\mathbb{R}^2)$ et A et B sont des opérateurs auto-adjoints qui presque commutent, alors

$$\big[\varphi(A,B),\psi(A,B)\big] = \iint\!\!\!\int \frac{\varphi(x,y_1)-\varphi(x,y_2)}{y_1-y_2}\,dE_A(x)\,dE_B(y_1)[B,\psi(A,B)]\,dE_B(y_2)$$
$$+ \iint\!\!\!\int \frac{\varphi(x_1,y)-\varphi(x_2,y)}{x_1-x_2}\,dE_A(x_1)[A,\psi(A,B)]\,dE_A(x_2)\,dE_B(y)$$

et

$$\big\|[\varphi(A,B),\psi(A,B)]\big\|_{\boldsymbol{S}_1} \le \text{const}\,\|\varphi\|_{B^1_{\infty,1}(\mathbb{R}^2)}\|\psi\|_{B^1_{\infty,1}(\mathbb{R}^2)}\big\|[A,B]\big\|_{\boldsymbol{S}_1}.$$

Ça implique que la formule des traces de Helton et Howe est vrai pour toutes les fonctions φ et ψ dans $B^1_{\infty,1}(\mathbb{R}^2)$.

---

1. **Introduction**

The spectral theorem allows one for a pair of commuting (bounded) self-adjoint operators A and B to construct a linear and multiplicative functional calculus

$$\varphi\mapsto\varphi(A,B)=\int_{\mathbb{R}^2}\varphi(x,y)\,dE_{A,B}(x,y)$$

for the class of bounded Borel functions on the plane $\mathbb{R}^2$. Here $E_{A,B}$ is the joint spectral measure of the pair(A,B) defined on the Borel subsets of $\mathbb{R}^2$.

If A and B are noncommuting self-adjoint operators, we can define functions of A and B in terms of double operator integrals

$$\varphi(A,B)\stackrel{\rm def}{=}\iint_{\mathbb{R}^2}\varphi(x,y)\,dE_A(x)\,dE_B(y) \tag{3}$$

for functions φ that are Schur multipliers with respect to the spectral measure E_A and E_B of the operators A and B. The theory of double operator integrals was developed by Birman and Solomyak [BS] (we also refer the reader to [Pe2] and [AP] for double operator integrals and Schur multipliers). It was observed in [ANP] the Besov space $B^1_{\infty,1}(\mathbb{R}^2)$ of functions on $\mathbb{R}^2$ is contained in the space of Schur multipliers with respect to compactly supported spectral measures on $\mathbb{R}$, and so for $\varphi\in B^1_{\infty,1}(\mathbb{R}^2)$, the operator $\varphi(A,B)$ is well defined by (3) for bounded self-adjoint operators A and B (see [Pee] for the definition and properties of Besov classes).

In this paper we deal with almost commuting self-adjoint operators. *Self-adjoint operators A and B are called almost commuting* if their *commutator* $[A,B]\stackrel{\rm def}{=}AB-BA$ belongs to trace class.

In [HH] the following trace formula was obtained for bounded almost commuting self-adjoint operators A and B:

$$\operatorname{trace}\big(\mathrm{i}\big(\varphi(A,B)\psi(A,B)-\psi(A,B)\varphi(A,B)\big)\big)=\iint_{\mathbb{R}^2}\left(\frac{\partial\varphi}{\partial x}\frac{\partial\psi}{\partial y}-\frac{\partial\varphi}{\partial y}\frac{\partial\psi}{\partial x}\right)dP, \tag{4}$$

where P is a signed Borel compactly supported measure that corresponds to the pair (A,B). The formula holds for polynomials φ and ψ.

It was shown in [Pin2] that the signed measure P is absolutely continuous with respect to planar Lebesgue measure and

$$dP(x,y)=\frac{1}{2\pi}g(x,y)\,dx\,dy,$$

where g is the *Pincus principal function*, which was introduced in [Pin1].

In [CP] the polynomial functional calculus for almost commuting self-adjoint operators was extended to a functional calculus for the class of functions $\varphi=\mathcal{F}\omega$ that are Fourier transforms of complex Borel measures ω on $\mathbb{R}^2$ satisfying

$$\int_{\mathbb{R}^2}(1+|t|)(1+|s|)\,d|\omega|(s,t)<\infty,$$

and the Helton–Howe trace formula (4) was extended to the class of such functions.

The problem of constructing a rich functional calculus, which would extend the functional calculus constructed in [CP] and for which trace formula (4) would still hold was considered in [Pe4]. The problem was to find a big class of functions $\mathcal{C}$ on $\mathbb{R}^2$ and construct a functional calculus $\varphi\mapsto\varphi(A,B)$, $\varphi\in\mathcal{C}$, that has the following properties:

(i) *the functional calculus $\varphi\mapsto\varphi(A,B)$, $\varphi\in\mathcal{C}$, is linear;*

(ii) *if $\varphi(s,t)=u(s)v(t)$, then $\varphi(A,B)=u(A)v(B)$;*

(iii) *if φ, $\psi\in\mathcal{C}$, then $\varphi(A,B)\psi(A,B)-\psi(A,B)\varphi(A,B)\in\boldsymbol{S}_1$;*

(iv) *formula* (4) *holds for arbitrary φ and ψ in $\mathcal{C}$.*

Note that the right-hand side of (4) makes sense for arbitrary Lipschitz functions φ and ψ. However, it was established in [Pe4] that a functional calculus satisfying (i) - (iii) cannot be defined for all continuously differentiable functions. This was deduced from the trace class criterion for Hankel operators (see [Pe1] and [Pe5]).

On the other hand, in [Pe4] estimates of [Pe2] and [Pe3] were used to construct a functional calculus satisfying (i) - (iv) for the class $\mathcal{C}=\big(L^\infty(\mathbb{R})\hat{\otimes}B^1_{\infty,1}(\mathbb{R})\big)\bigcap\big(B^1_{\infty,1}(\mathbb{R})\hat{\otimes}L^\infty(\mathbb{R})\big)$. Here $\hat{\otimes}$ stands for projective tensor product and $B^1_{\infty,1}(\mathbb{R})$ is a Besov class.

In this paper we considerably enlarge the class of functions $\big(L^\infty(\mathbb{R})\hat{\otimes}B^1_{\infty,1}(\mathbb{R})\big)\bigcap\big(B^1_{\infty,1}(\mathbb{R})\hat{\otimes}L^\infty(\mathbb{R})\big)$ and construct a functional calculus satisfying (i) - (iv) for the Besov class $B^1_{\infty,1}(\mathbb{R}^2)$ of functions of two variables.

2. **Triple operator integrals**

Triple operator integrals

$$\iiint \Phi(x_1,x_2,x_3)\,dE_1(x_1)T\,dE_2(x_2)R\,dE_3(x_3) \tag{5}$$

were defined in [Pe6] for spectral measures E_1, E_2, E_3, bounded linear operators T and R, and for functions Φ in the integral projective tensor product $L^\infty(E_1)\hat{\otimes}_{\rm i}L^\infty(E_2)\hat{\otimes}_{\rm i}L^\infty(E_3)$.

Later the definition of triple operator integrals was extended in [JTT] to the class of functions Φ in the Haagerup tensor product $L^\infty(E_1)\otimes_{\rm h}L^\infty(E_2)\otimes_{\rm h}L^\infty(E_3)$ (see [Pis] for imformation about Haagerup tensor products). However, it was shown in [ANP] that unlike in the case $\Phi\in L^\infty(E_1)\hat{\otimes}_{\rm i}L^\infty(E_2)\hat{\otimes}_{\rm i}L^\infty(E_3)$, the condition that Φ belongs to $L^\infty(E_1)\otimes_{\rm h}L^\infty(E_2)\otimes_{\rm h}L^\infty(E_3)$ does not guarantee that if one of the operators T and R is of trace class, then the triple operator integral (5) belongs to $\boldsymbol{S}_1$.

In [ANP] the following Haagerup-like tensor products were introduced:

Definition. *A function Ψ is said to belong to the tensor product $L^\infty(E_1)\otimes_{\rm h}L^\infty(E_2)\otimes^{\rm h}L^\infty(E_3)$ if it admits a representation*

$$\Psi(x_1,x_2,x_3)=\sum_{j,k\ge0}\alpha_j(x_1)\beta_k(x_2)\gamma_{jk}(x_3) \tag{6}$$

with $\{\alpha_j\}_{j\ge0},\ \{\beta_k\}_{k\ge0}\in L^\infty(\ell^2)$ and $\{\gamma_{jk}\}_{j,k\ge0}\in L^\infty(\mathcal{B})$, where $\mathcal{B}$ is the space of bounded operators on ℓ^2.

For a bounded linear operator R and for a trace class operator T, the triple operator integral

$$W=\iiint \Psi(x_1,x_2,x_3)\,dE_1(x_1)T\,dE_2(x_2)R\,dE_3(x_3)$$

was defined in [ANP] as the following continuous linear functional on the class of compact operators:

$$Q\mapsto\operatorname{trace}\left(\left(\iiint\Psi(x_1,x_2,x_3)\,dE_2(x_2)R\,dE_3(x_3)Q\,dE_1(x_1)\right)T\right) \tag{7}$$

and it was shown that

$$\|W\|_{\boldsymbol{S}_1}\le\|\Psi\|_{L^\infty\otimes_{\rm h}L^\infty\otimes^{\rm h}L^\infty}\|T\|_{\boldsymbol{S}_1}\|R\|,$$

where $\|\Psi\|_{L^\infty\otimes_{\rm h}L^\infty\otimes^{\rm h}L^\infty}$ is the infimum of $\|\{\alpha_j\}_{j\ge0}\|_{L^\infty(\ell^2)}\|\{\beta_k\}_{k\ge0}\|_{L^\infty(\ell^2)}\|\{\gamma_{jk}\}_{j,k\ge0}\|_{L^\infty(\mathcal{B})}$ over all representations in (6).

Similarly, the tensor product $L^\infty(E_1)\otimes^{\rm h}L^\infty(E_2)\otimes_{\rm h}L^\infty(E_3)$ was defined in [ANP] as the class of functions Ψ of the form

$$\Psi(x_1,x_2,x_3)=\sum_{j,k\ge0}\alpha_{jk}(x_1)\beta_j(x_2)\gamma_k(x_3)$$

where $\{\beta_j\}_{j\ge0},\ \{\gamma_k\}_{k\ge0}\in L^\infty(\ell^2)$, $\{\alpha_{jk}\}_{j,k\ge0}\in L^\infty(\mathcal{B})$. If T is a bounded linear operator, and $R\in\boldsymbol{S}_1$, then the continuous linear functional

$$Q\mapsto\operatorname{trace}\left(\left(\iiint\Psi(x_1,x_2,x_3)\,dE_3(x_3)Q\,dE_1(x_1)T\,dE_2(x_2)\right)R\right)$$

on the class of compact operators determines a trace class operator

$$W\stackrel{\rm def}{=}\iiint\Psi(x_1,x_2,x_3)\,dE_1(x_1)T\,dE_2(x_2)R\,dE_3(x_3)$$

and

$$\|W\|_{\boldsymbol{S}_1}\le\|\Psi\|_{L^\infty\otimes^{\rm h}L^\infty\otimes_{\rm h}L^\infty}\|T\|\cdot\|R\|_{\boldsymbol{S}_1}.$$

3. **Commutators of functions of almost commuting self-adjoint operators**

Given a differentiable function φ on $\mathbb{R}^2$, we define the divided differences $\mathfrak{D}_1 f$ and $\mathfrak{D}_2 f$ on $\mathbb{R}^3$ by

$$(\mathfrak{D}_1\varphi)(x_1,x_2,y) \stackrel{\text{def}}{=} \frac{\varphi(x_1,y)-\varphi(x_2,y)}{x_1-x_2} \quad \text{and} \quad (\mathfrak{D}_2\varphi)(x,y_1,y_2) = \frac{\varphi(x,y_1)-\varphi(x,y_2)}{y_1-y_2}.$$

It was shown in [ANP] that if φ is a bounded function on $\mathbb{R}^2$ whose Fourier transform is supported in the ball $\{\xi \in \mathbb{R}^2 : \ \|\xi\| \le 1\}$, then

$$(\mathfrak{D}_1\varphi)(x_1,x_2,y) = \sum_{j,k\in\mathbb{Z}} \frac{\sin(x_1-j\pi)}{x_1-j\pi}\cdot\frac{\sin(x_2-k\pi)}{x_2-k\pi}\cdot\frac{\varphi(j\pi,y)-\varphi(k\pi,y)}{j\pi-k\pi}.$$

$$\sum_{j\in\mathbb{Z}} \frac{\sin^2(x_1-j\pi)}{(x_1-j\pi)^2} = \sum_{k\in\mathbb{Z}} \frac{\sin^2(x_2-k\pi)}{(x_2-k\pi)^2} = 1, \quad x_1\ x_2 \in \mathbb{R},$$

and

$$\sup_{y\in\mathbb{R}} \left\| \left\{ \frac{\varphi(j\pi,y)-\varphi(k\pi,y)}{j\pi-k\pi} \right\}_{j,k\in\mathbb{Z}} \right\|_{\mathcal{B}} \le \text{const}\, \|f\|_{L^\infty(\mathbb{R})}.$$

It follows that for $\varphi \in B^1_{\infty,1}(\mathbb{R}^2)$,

$$\mathfrak{D}_1\varphi \in L^\infty(\mathbb{R}) \otimes_{\rm h} L^\infty(\mathbb{R}) \otimes^{\rm h} L^\infty(\mathbb{R}) \quad \text{and} \quad \|\varphi\|_{L^\infty(\mathbb{R})\otimes_{\rm h}L^\infty(\mathbb{R})\otimes^{\rm h}L^\infty(\mathbb{R})} \le \text{const}\, \|\varphi\|_{B^1_{\infty,1}}.$$

Similarly, for $\varphi \in B^1_{\infty,1}(\mathbb{R}^2)$,

$$\mathfrak{D}_2\varphi \in L^\infty(\mathbb{R}) \otimes^{\rm h} L^\infty(\mathbb{R}) \otimes_{\rm h} L^\infty(\mathbb{R}) \quad \text{and} \quad \|\varphi\|_{L^\infty(\mathbb{R})\otimes^{\rm h}L^\infty(\mathbb{R})\otimes_{\rm h}L^\infty(\mathbb{R})} \le \text{const}\, \|\varphi\|_{B^1_{\infty,1}}.$$

Theorem 3.1. *Let A and B be self-adjoint operators and let Q be a bounded linear operator such that $[A,Q] \in \boldsymbol{S}_1$ and $[B,Q] \in \boldsymbol{S}_1$. Suppose that $\varphi \in B^1_{\infty,1}(\mathbb{R}^2)$. Then $\big[\varphi(A,B),Q\big] \in \boldsymbol{S}_1$,*

$$\begin{aligned}\big[\varphi(A,B),Q\big] &= \iiint \frac{\varphi(x,y_1)-\varphi(x,y_2)}{y_1-y_2}\,dE_A(x)\,dE_B(y_1)[B,Q]\,dE_B(y_2)\\ &\quad+\iiint \frac{\varphi(x_1,y)-\varphi(x_2,y)}{x_1-x_2}\,dE_A(x_1)[A,Q]\,dE_A(x_2)\,dE_B(y)\end{aligned} \tag{8}$$

and

$$\big\|[\varphi(A,B),Q]\big\|_{\boldsymbol{S}_1} \le \text{const}\, \|\varphi\|_{B^1_{\infty,1}(\mathbb{R}^2)}\big(\big\|[A,Q]\big\|_{\boldsymbol{S}_1} + \big\|[B,Q]\big\|_{\boldsymbol{S}_1}\big).$$

To obtain the main result of the paper, we apply Theorem 3.1 in the case $Q = \psi(A,B)$, where $\psi \in B^1_{\infty,1}(\mathbb{R}^2)$.

Theorem 3.2. *Let A and B be almost commuting self-adjoint operators and let φ and ψ be functions in the Besov class $B^1_{\infty,1}(\mathbb{R}^2)$. Then*

$$\begin{aligned}\big[\varphi(A,B),\psi(A,B)\big] &= \iiint \frac{\varphi(x,y_1)-\varphi(x,y_2)}{y_1-y_2}\,dE_A(x)\,dE_B(y_1)[B,\psi(A,B)]\,dE_B(y_2)\\ &\quad+\iiint \frac{\varphi(x_1,y)-\varphi(x_2,y)}{x_1-x_2}\,dE_A(x_1)[A,\psi(A,B)]\,dE_A(x_2)\,dE_B(y)\end{aligned} \tag{9}$$

and

$$\big\|[\varphi(A,B),\psi(A,B)]\big\|_{\boldsymbol{S}_1} \le \text{const}\, \|\varphi\|_{B^1_{\infty,1}(\mathbb{R}^2)}\|\psi\|_{B^1_{\infty,1}(\mathbb{R}^2)}\big\|[A,B]\big\|_{\boldsymbol{S}_1}. \tag{10}$$

Note that the right-hand side of inequality (10) does not involve the norms of A or B. Thus formulae (8) and (9) allow us to consider commutators $\big[f(A,B),g(A,B)\big]$ even for unbounded self-adjoint operators A and B with trace class commutator $[A,B]$, though the the functions $f(A,B)$ and $g(A,B)$ of A and B are not necessarily defined for f and g in $B^1_{\infty,1}(\mathbb{R}^2)$.

Let us also mention that in [Pe4] it was proved that for almost commuting self-adjoint operators A and B, the functional calculus $\varphi\mapsto\varphi(A,B)$, $\varphi\in\big(L^\infty(\mathbb{R})\hat{\otimes}B^1_{\infty,1}(\mathbb{R})\big)\bigcap\big(B^1_{\infty,1}(\mathbb{R})\hat{\otimes}L^\infty(\mathbb{R})\big)$, is *almost multiplicative*, i.e.,

$$(\varphi\psi)(A,B)-\varphi(A,B)\psi(A,B)\in\boldsymbol{S}_1,\quad\varphi,\psi\in\big(L^\infty(\mathbb{R})\hat{\otimes}B^1_{\infty,1}(\mathbb{R})\big)\bigcap\big(B^1_{\infty,1}(\mathbb{R})\hat{\otimes}L^\infty(\mathbb{R})\big).$$

It would be interesting to find out whether the functional calculus $\varphi\mapsto\varphi(A,B)$, $\varphi\in B^1_{\infty,1}(\mathbb{R}^2)\bigcap L^\infty(\mathbb{R}^2)$, is also almost multiplicative.

4. **An extension of the Helton–Howe trace formula**

In this section we use the results of the previous section to extend the Helton–Howe trace formula.

Theorem 4.1. *Let A and B be almost commuting self-adjoint operators and let φ and ψ be functions in the Besov class $B^1_{\infty,1}(\mathbb{R}^2)$. Then the following formula holds:*

$$\operatorname{trace}\big(\mathrm{i}\big(\varphi(A,B)\psi(A,B)-\psi(A,B)\varphi(A,B)\big)\big)=\frac{1}{2\pi}\iint_{\mathbb{R}^2}\left(\frac{\partial\varphi}{\partial x}\frac{\partial\psi}{\partial y}-\frac{\partial\varphi}{\partial y}\frac{\partial\psi}{\partial x}\right)g(x,y)\,dx\,dy,\tag{11}$$

where g is the Pincus principal function associated with the operators A and B.

It would be interesting to extend the Pincus principal function to the case of unbounded self-adjoint operators with trace class commutators and extend formula (11) to unbounded almost commuting operators.

References

[ANP] A.B. Aleksandrov, F.L. Nazarov and V.V. Peller, *Functions of perturbed noncommuting self-adjoint operators*, to appear in C.R Acad. Sci Paris

[AP] A.B. Aleksandrov and V.V. Peller, *Operator Hölder–Zygmund functions*, Advances in Math. **224** (2010), 910-966.

[BS] M.S. Birman and M.Z. Solomyak, *Double Stieltjes operator integrals*, Problems of Math. Phys., Leningrad. Univ. **1** (1966), 33–67 (Russian). English transl.: Topics Math. Physics **1** (1967), 25–54, Consultants Bureau Plenum Publishing Corporation, New York.

[CP] R.W. Carey and J.D. Pincus, *Mosaics, principal functions and mean motion in von Neumann algebras*, Acta Math. **138** (1977), 153–218.

[Cl] K.F. Clancey, *Seminormal operators*, Lecture Notes in Math. **742**, Springer–Verlag, Berlin, 1979.

[HH] J.W. Helton and R. Howe, *Integral operators, commutators, traces, index, and homology*, in "Lecture Notes in Math.", vol. **345**, pp. 141–209, Springer-Verlag, New York, 1973.

[JTT] K. Juschenko, I.G. Todorov and L. Turowska, *Multidimensional operator multipliers*, Trans. Amer. Math. Soc. **361** (2009), 4683-4720.

[Pee] J. Peetre, *New thoughts on Besov spaces*, Duke Univ. Press., Durham, NC, 1976.

[Pe1] V.V. Peller, *Hankel operators of class $\mathbf{S}_p$ and their applications (rational approximation, Gaussian processes, the problem of majorizing operators)*, Mat. Sbornik, **113** (1980), 538-581. English Transl. in Math. USSR Sbornik, **41** (1982), 443-479.

[Pe2] V.V. Peller, *Hankel operators in the theory of perturbations of unitary and self-adjoint operators*, Funktsional. Anal. i Prilozhen. **19:2** (1985), 37–51 (Russian). English transl.: Funct. Anal. Appl. **19** (1985) , 111–123.

[Pe3] V.V. Peller, *Hankel operators in the perturbation theory of of unbounded self-adjoint operators.* Analysis and partial differential equations, 529–544, Lecture Notes in Pure and Appl. Math., **122**, Dekker, New York, 1990.

[Pe4] V.V. Peller, *Functional calculus for a pair of almost commuting self-adjoint operators*, J. Funct. Anal. **112** (1993), 325-345.

[Pe5] V.V. Peller, *Hankel operators and their applications,* Springer-Verlag, New York, 2003.

[Pe6] V.V. Peller, *Multiple operator integrals and higher operator derivatives*, J. Funct. Anal. **233** (2006), 515–544.

[Pin1] J.D. Pincus, *Commutators and systems of singular integral equations, I*, Acta Math. **121** (1968), 219-249.

[Pin2] J.D. Pincus, *On the trace of commutators in the algebra of operators generated by an operator with trace class self-commutator*, Stony Brook preprint (1972).

[Pis] G. Pisier, *Introduction to operator space theory*, London Math. Society Lect. Notes series **294**, Cambridge University Press, 2003.